



HARNESSING MICROBIAL CONSORTIA FOR NEXT-GENERATION BIOFERTILIZERS AND BIOCONTROL AGENTS: MECHANISMS, CHALLENGES, AND OPPORTUNITIES. A REVIEW

¹Ayuba, K. S., ²Ogbonnaya, C. S., ³Tanko, F. and ⁴Yahaya, M.

¹Department of Biological Sciences, Abubakar Tafawa Balewa University, Bauchi

²Department of Soil Science, Abubakar Tafawa Balewa University, Bauchi

³Department of Soil Science, Federal University, Minna

⁴College of Agriculture Kabba, Division of Agricultural Colleges, Ahmadu Bello University, Zaria

Corresponding Author's Email: shiayetinfo@gmail.com,

ABSTRACT

Nigerian agriculture faces persistent soil fertility depletion, rising costs of synthetic fertilizers, and environmental degradation, necessitating the need for sustainable, environmentally friendly and cost-effective alternatives to chemical inputs. This review evaluates the application of microbial consortia as biofertilizers and biocontrol agents in Nigeria, focusing on their benefits, constraints, and prospects for adoption. Relevant empirical studies on plant growth-promoting microorganisms, multi-strain inoculants, and biological pest suppression were reviewed to assess agronomic performance, implementation challenges, emerging innovations and adoption constraints. Microbial consortia enhance biological nitrogen fixation, phosphorus solubilization, nutrient cycling, plant defence responses, and phytohormone production leading to improved crop growth, soil health, and resilience to abiotic stress. Beneficial taxa such as *Trichoderma*, *arbuscular mycorrhizal fungi* and *Pseudomonas* suppress soil-borne pathogens, reducing pesticide dependence. However, inconsistent field performance, formulation challenges, short shelf life, weak quality control, limited farmer awareness, and inadequate production and regulatory systems constrain large-scale adoption. With improved formulation technologies, development of indigenous microbial strains, stronger extension services, and supportive policies, microbial consortia offer a viable pathway toward sustainable intensification, reduced agrochemical reliance, and enhanced productivity in Nigerian agriculture.

Keywords: Microbial-consortia, Biofertilizer, Biocontrol, Sustainable, Agriculture.

INTRODUCTION

Growing environmental concerns, declining soil fertility, and increasing regulatory restrictions on synthetic fertilizers and pesticides have intensified global interest in sustainable, microbe-based agricultural inputs (Saad *et al.*, 2020; Shahwar *et al.*, 2023). Excessive agrochemical use has contributed to soil degradation, groundwater contamination, biodiversity loss, and greenhouse gas emissions, thereby threatening long-term productivity and food security (Tilman *et al.*, 2011; Pretty *et al.*, 2018). Consequently, biofertilizers and biocontrol agents derived from beneficial microorganisms are gaining recognition as environmentally sound alternatives capable of enhancing crop performance while preserving ecosystem integrity. Conventional biofertilizer strategies have primarily relied on single-strain inoculants, including nitrogen-fixing bacteria (*Rhizobium*, *Azotobacter*), phosphate-solubilizing microorganisms (*Bacillus*, *Pseudomonas*), and arbuscular mycorrhizal fungi (Asif *et al.*, 2018). Although these inputs show promising results under controlled conditions, their effectiveness in field environments is often inconsistent due to variations in soil physical-chemical properties, climatic fluctuations, and competition with indigenous microbiota (Herrmann & Lesueur, 2013; Malusá *et al.*, 2016). Such variability underscores the limitations of single-strain approaches in addressing the complexity of agroecosystems. Agricultural soils are dynamic and biologically diverse systems where plant growth depends on interactions among multiple microbial taxa, plant roots, and abiotic factors (Fierer, 2017). In natural environments, microorganisms operate within cooperative networks that collectively regulate nutrient cycling, plant growth promotion, and disease suppression (Philippot *et al.*, 2013). Hence, individual strains rarely replicate the functional resilience of native soil microbial communities. In response, microbial consortia: structured assemblages of interacting bacteria, fungi,

and other beneficial microorganisms are increasingly viewed as next-generation biofertilizers and biocontrol agents (Nunes *et al.*, 2024; Singh *et al.*, 2024). Figure 1 shows the potential contribution of microbial consortia to sustainable agriculture.




Figure 1: *Microbial Consortia Contribute to Sustainable and Resilient Agriculture*

Source: Authors' Modification (Seenivasagan & Babalola, 2021; Rivas-García *et al.*, 2025).

Through synergistic and complementary interactions, these consortia simultaneously perform multiple plant-beneficial functions, including biological nitrogen fixation, nutrient solubilization, phytohormone production, and pathogen suppression (Bashan *et al.*, 2014; Trivedi *et al.*, 2020). Compared with single-strain inoculants, they exhibit greater ecological stability, functional redundancy, and adaptability to environmental stress, resulting in improved persistence and performance under field conditions (Niu *et al.*, 2017; de Souza *et al.*, 2020). Therefore, understanding of microbial consortia and their functional potential is essential for advancing sustainable agricultural systems.

Defining Microbial Consortia

Microbial consortia are structured communities of two or more interacting microorganisms that cooperate to deliver ecological or agronomic benefits superior to those of single strains. In agricultural systems, these consortia typically comprise bacteria, fungi, and actinomycetes inhabiting the rhizosphere or endosphere, where coordinated metabolic activities enhance nutrient acquisition, plant growth, and disease suppression (Woo & Pepe, 2018; Santoyo *et al.*, 2021). Unlike simple inoculant mixtures, microbial consortia are defined by functional complementarity and interspecies interactions such as cross-feeding, quorum sensing, and syntrophic metabolism. These interactions generate synergistic or emergent functions, for example, simultaneous nitrogen fixation, phosphorus solubilization, phytohormone production, and antimicrobial activity resulting in multifunctional and more stable performance under variable field conditions (Bashan *et al.*, 2014; Santoyo *et al.*, 2021). Natural soil microbiomes inherently function as complex consortia shaped by root exudates and environmental drivers. Advances in microbial ecology and systems biology now enable the deliberate design of synthetic microbial communities (SynComs) that replicate these natural associations, improving reliability, resilience, and broad-spectrum crop benefits (Toju *et al.*, 2018; Woo & Pepe, 2018). Overall, microbial consortia can be viewed as integrated, multispecies systems selected or engineered to synergistically improve soil fertility, crop productivity, and plant health, forming the foundation of next-generation biofertilizers and biocontrol agents.

MECHANISMS OF ACTION

Microbial consortia operate through coordinated, multi-layered biochemical and ecological processes within the rhizosphere. Unlike single-strain inoculants that perform isolated functions, multispecies consortia combine complementary metabolic traits to simultaneously enhance nutrient mobilization, regulate plant growth, and suppress pathogens. These synergistic interactions result in more stable and consistent crop responses across heterogeneous soil environments. The principal mechanisms include nutrient acquisition and cycling, phytohormone modulation, and biological disease control.

Nutrient Acquisition and Cycling

Soil fertility constraints often arise from limited nutrient bioavailability rather than total nutrient abundance. Although nitrogen, phosphorus, and potassium may be present in soils, they frequently occur in chemically inaccessible forms. Microbial consortia overcome these constraints through complementary enzymatic transformations that convert unavailable nutrient pools into plant-assimilable forms, often described by “*the division of labour model*” (Qian *et al.*, 2020). Microbial consortia also drive the cycling of elements through interlinked biochemical pathways. Such as shared metabolic intermediates; for instance, one microbe may consume the waste product of another, facilitating the complete decomposition of complex organic compounds. Microbes communicate through chemical signals (e.g., quorum sensing), which coordinate metabolic activities like enzyme production and antibiotic synthesis to modulate nutrient availability (de Jesús Astacio *et al.*, 2021). Rhizosphere interactions between roots and beneficial microbes that underpin these processes are illustrated in Figure 2, highlights symbiotic colonization and nutrient exchange at the soil–root interface.




Figure 2. Rhizosphere Processes and Symbiotic Plant–microbe Interactions Supporting Nutrient Mobilization.

A: Some rhizosphere processes in the soil B: Simplified diagram of the relation between the plant and the symbiotic bacteria (cyan) in the root nodules

Source: Wikipedia (2026)

Biological Nitrogen Fixation (BNF)

Biological nitrogen fixation is a key ecosystem service provided by diazotrophic members of microbial consortia, including *Rhizobium*, *Bradyrhizobium*, *Azospirillum*, and *Azotobacter*. These microorganisms reduce atmospheric N₂ to ammonia via nitrogenase providing sustainable plant nutrition, particularly within protected microaerobic niches such as legume root nodules, thereby enabling direct nitrogen assimilation by plants (Herridge *et al.*, 2008). In consortia, cross-feeding and

biofilm formation enhance energy supply, persistence, and root colonization efficiency, thereby increasing nitrogen contribution relative to single-strain inoculants. These synergistic, multi-species teams enhance nitrogen-fixation efficiency, improve soil health, and reduce reliance on synthetic fertilizers, offering a greener agricultural alternative (Dey *et al.*, 2021; Herrero *et al.*, 2025).

Phosphate Solubilization and Mineralization

Despite high total phosphorus content, most soil phosphorus in Nigeria remains immobilized in insoluble complexes. Phosphate-solubilizing microorganisms such as *Bacillus*, *Pseudomonas*, *Aspergillus*, and *Penicillium* mobilize these reserves through organic acid secretion, rhizosphere acidification, chelation of metal ions, and phosphatase activity, releasing plant-available orthophosphate (Prasad & Chakraborty, 2019). Within consortia, fungal hyphae further extend nutrient capture zones, enhancing phosphorus uptake efficiency in low p soils. The transformations and pathways governing these dynamics are summarized in Figure 3.




Figure 3: Soil Phosphorus Cycle: Showing the Sources of Phosphorus Inputs in the Soil, Pathways through which Phosphorus becomes Available/unavailable for Plant Uptake and Phosphorus Outputs/loss Pathways.

Source: Prasad and Chakraborty (2019)

Potassium and Micronutrient Mobilization

Potassium and micronutrients such as Fe, Zn, and Mn are commonly locked within mineral matrices or oxidized forms. Certain *Bacillus* and *Frateria* species release organic acids and exopolysaccharides that weather minerals and liberate exchangeable K⁺. Concurrently, siderophore-producing microbes chelate Fe³⁺, enhancing plant uptake while restricting pathogen access, thereby contributing to both nutrition and disease suppression (Zhang & Smith, 2020).

PHYTOHORMONE MODULATION

Beyond nutrient mobilization, microbial consortia directly regulate plant physiology through the synthesis and modulation of phytohormones that shape root architecture, shoot development, and stress responses (Egamberdieva *et al.*, 2017; Kunal *et al.*, 2023). These microorganisms produce or modulate hormones such as auxins, cytokinins, gibberellins, and ethylene-related compounds, thereby reshaping root system architecture and enhancing nutrient acquisition efficiency (Egamberdieva *et al.*, 2017; Jia *et al.*, 2023). By altering hormonal balance in the rhizosphere, these microorganisms indirectly enhance adaptive responses to environmental stress overall plant vigour (Jia *et al.*, 2023; Feng *et al.*, 2025). The spatial coordination of hormone-mediated root responses and microbial signalling is illustrated in Figure 4, which highlights auxin-driven root hair development under nutrient limitation.




Figure 4: Auxin-mediated Epidermal Signalling and Root Hair Proliferation under Nutrient Stress
Source: Jia *et al.* (2023).

Auxin (Indole-3-Acetic Acid, IAA) is the most widely reported microbial phytohormone (Spaepen & Vanderleyden, 2011; Jia *et al.*, 2023; Etesami & Glick, 2024). Produced by genera such as *Pseudomonas*, *Azospirillum*, and *Bacillus*, indole-3-acetic acid (IAA) stimulates lateral root formation, root hair proliferation, and rhizosphere exploration, thereby enlarging the absorptive surface area for water and nutrients (Spaepen & Vanderleyden, 2011). In microbial consortia, cumulative auxin production often generates stronger and more consistent growth responses than single strains (Kumar *et al.*, 2024; Meza *et al.*, 2025). Meza *et al.* (2025) reported auxin-producing PGPB (*Bacillus proteolyticus* Cyn1 and *Bacillus safensis* Cyn2) found their consortium produced 90.67 µg/mL IAA, exceeding individual strains (46 µg/mL and 43.78 µg/mL), due to synergistic effects that support stronger growth promotion. Several rhizobacteria synthesize cytokinins and gibberellins or influence endogenous plant pathways. Cytokinins promote cell division and delay senescence, whereas gibberellins enhance stem elongation, seed germination, and biomass accumulation, collectively improving canopy development and yield potential.

BIOCONTROL AND DISEASE SUPPRESSION

Microbial consortia function as effective biological control agents by protecting plants against soil-borne and foliar pathogens through coordinated direct antagonism and host-mediated defence activation. Their suppressive capacity arises from multiple complementary mechanisms that operate simultaneously in the rhizosphere. Direct antagonism occurs through antibiosis production and lytic enzyme activity. Beneficial rhizobacteria produce antimicrobial metabolites such as phenazines, pyrrolnitrin, hydrogen cyanide, and cyclic lipopeptides (surfactin, iturin, and fengycin), which inhibit spore germination, disrupt pathogen membranes, and suppress growth (Saeed *et al.*, 2021; Dutta, 2022). Fungal partners, particularly *Trichoderma* spp., secrete hydrolytic enzymes including chitinases and glucanases that degrade fungal cell walls, thereby enhancing pathogen lysis (Khatri *et al.*, 2017). The combine action of bacterial antibiotics and fungal enzymatic degradation results in synergistic disease suppression.

Resource competition and niche exclusion further limit pathogen establishment. Rapid rhizosphere colonization, mediated by bacterial motility and chemotaxis, enables beneficial microbes to monopolize nutrients and attachment sites (Liu *et al.*, 2024). By depleting available carbon sources and sequestering iron through siderophore production, consortia create nutrient-limited conditions unfavourable to invading pathogens. Additionally, biofilm formation on the rhizoplane stabilizes microbial communities, improves persistence, and restricts pathogen access. In *Pseudomonas fluorescens*, robust biofilm formation is strongly associated with suppression of rhizosphere pathogens (Liu *et al.*, 2024a). Disease reduction against soilborne *Ralstonia solanacearum* following consortium inoculation further highlights the effectiveness of competitive exclusion (Gu *et al.*, 2020). Beyond direct

suppression and competitive exclusion microbial consortia activate plant immune responses through induced systemic resistance (ISR). ISR is mediated primarily via jasmonic acid and ethylene signalling pathways, which prime plants for faster and stronger defence responses upon pathogen challenge. This primed state enhances protection without the metabolic costs of continuous defence activation, thereby improving overall plant fitness (Nunes *et al.*, 2024; Salwan *et al.*, 2023). The ISR signalling cascade and its role in systemic protection are illustrated in Fig. 5. Collectively, these mechanisms: antibiosis, enzymatic lysis, competitive exclusion, biofilm establishment, and ISR act synergistically, enabling microbial consortia to provide reliable and sustainable disease suppression in agricultural systems.




Figure 5: Induced Systemic Resistance (ISR) Signalling in Plants.
Source: Kumari *et al.* (2024a).

CONSORTIUM FORMULATION AND DELIVERY

The agronomic performance of microbial consortia depends not only on the functional traits of constituent strains but also on their stability, compatibility, and successful delivery to the rhizosphere (Singh *et al.*, 2025). Although laboratory assays frequently demonstrate strong plant growth-promoting or biocontrol potential, field efficacy is often inconsistent due to poor survival during storage, environmental stress after application, and inadequate root colonization. Consortium design typically follows three interrelated principles: functional complementarity, ecological compatibility, and environmental resilience (Keerthana *et al.*, 2024; Singh *et al.*, 2025). Table 1 shows examples of consortium formulation and their application

Table 1: Examples of Consortium Formulations and their Applications

Formulation Type	Microbial Composition	Application / Observed Effects	Reference
Kaolinite, Bentonite, Lignite, Liquid & Na-Alginate Formulations	Consortia of N-fixers (Azospirillum or Gluconacetobacter), P-solubilizer (<i>Pseudomonas striata</i>) and K-solubilizer (KSB)	Enhanced N, P, K content in maize; liquid and alginate carriers improved nutrient content and shelf life.	Ashwini (2023).
Wettable Powder (kaolinite / diatomaceous earth) & Liquid	<i>The Great Five</i> bacterial consortium (A294, A167, H145, H440, H469)	Improved shelf stability and formulation performance for soft-rot control in potato storage.	Maciag <i>et al.</i> (2020).
Synthetic Consortia (SynCom)	Defined microbial communities (e.g., <i>Bacillus</i> sp. with AM fungi)	Enhanced plant growth, nutrient cycling, stress resilience and better adaptation to soil environment.	Singh <i>et al.</i> (2025).
Synergistic Microbial Consortia for Biocontrol	Selected rhizobacteria and fungi screened against <i>Fusarium oxysporum</i>	Enhanced systemic resistance in cumin against <i>Fusarium</i> wilt.	Singh <i>et al.</i> (2025a)
Trichoderma + Pseudomonas Consortium	<i>Trichoderma harzianum</i> + <i>T. asperellum</i> + <i>Pseudomonas fluorescens</i>	Suppressed bacterial blight in basmati rice and improved disease control.	Teja <i>et al.</i> (2025).
PGPR Rhizosphere Consortium	Multiple PGPR (e.g., <i>Bradyrhizobium</i> , <i>Pseudomonas</i> , <i>Bacillus</i> , nutrient solubilizers)	Promoted plant growth via multiple nutrient-mobilizing and PGP traits	Keerthana <i>et al.</i> (2024)


The effectiveness of these is strongly influenced by how efficiently the microorganisms are delivered, protected, and established in the rhizosphere. Despite demonstrated synergistic functions field performance often declines due to desiccation, temperature stress, UV exposure, and competition with native microbiota. Consequently, delivery systems must maintain cell viability during storage, preserve compatibility among consortium members, and enable controlled release at the root–soil interface (Bashan *et al.*, 2014; Malusá & Vassilev, 2014). Liquid inoculants remain the simplest and most widely adopted delivery method due to ease of production, handling, and compatibility with mechanized sowing (Chai *et al.*, 2022). Their application is often through seed coating, root dipping, or fertigation, ensuring rapid activation and uniform distribution. However, limited shelf life and sensitivity to environmental conditions may reduce microbial survival (Herrmann & Lesueur 2013). In contrast, solid and granular carriers such as peat, talc, lignite, compost, or biochar provide greater physical protection and gradual microbial release, enhancing persistence in soil. Biochar-based carriers are particularly advantageous due to their high porosity, which promotes microbial attachment while simultaneously improving soil moisture retention and nutrient availability (Chargrow, 2019; Gunal, 2025). Encapsulation technologies is another method that represent an advanced delivery strategy. Microbial cells are immobilized in polymeric matrices (e.g., alginate or chitosan beads), forming protective microenvironments that reduce desiccation and UV damage while enabling sustained release near plant roots. Encapsulation also minimizes antagonistic interactions among strains during storage, thereby improving the stability and consistency of multi-strain formulations (Maciag *et al.*, 2020; Kumar & Mohideen, 2025). The principal delivery approaches; liquid application, granular/solid carriers, and encapsulated systems are illustrated in Figure 5 – 7.




Figure 5: Root-dip inoculation of seedlings to enhance immediate root colonization
Source: Repáč *et al.* (2017).



Figure 6: Close-up view of pelletized biochar
Source: Page-Dumroese *et al.* (2016)



(a)



(b)

Figure 7: (a) Photographs of alginate gel beads and (b) photographs of alginate capsules.

EVIDENCE FROM NIGERIAN STUDIES

Empirical evidence from Nigeria consistently shows that microbial inoculants and consortia enhance crop productivity and soil fertility, particularly under low-input systems. Field studies report that biofertilizers improve nutrient availability, biological nitrogen fixation, and soil health, thereby reducing dependence on mineral fertilizers (Taiwo *et al.*, 2018; Bebeley *et al.*, 2024). Legume–rhizobia symbioses remain the most documented success. Inoculation of soybean, cowpea, and groundnut with compatible *Rhizobium* strains significantly increases nodulation, nitrogen fixation, and grain yield in nitrogen-deficient soils, often reducing or eliminating synthetic nitrogen use. Yield gains of 20 – 60 % have been reported, confirming rhizobia as a cost-effective strategy for smallholder farmers (Sanginga *et al.*, 2002; Abdulkadir *et al.*, 2021). Beyond legumes, plant growth–promoting rhizobacteria (PGPR) such as *Azospirillum*, *Bacillus*, and *Pseudomonas* enhance root development, phosphorus solubilization, and phytohormone production, leading to greater biomass and grain yield in cereals including maize and sorghum (Olanrewaju *et al.*, 2017). Multi-strain inoculation generally outperforms single isolates due to functional complementarity, integrating nutrient mobilization, nitrogen fixation, and stress tolerance (Adesemoye & Egamberdieva, 2013). Arbuscular mycorrhizal fungi (AMF) also show strong benefits in degraded tropical soils. By improving phosphorus uptake, drought tolerance, and nutrient acquisition, AMF enhance productivity of crops such as cassava and maize under reduced fertilizer inputs. These benefits are particularly valuable in the semi-arid regions of northern Nigeria where soil organic matter and available phosphorus are low (Bello *et al.*, 2024).

Recent research increasingly emphasizes consortium-based formulations that integrate rhizobia, PGPR, and fungi to deliver multiple functions simultaneously. These consortia improve soil



organic carbon, microbial biomass, and nutrient cycling while maintaining yields comparable to, or exceeding, chemical fertilizer systems (Agbodjato & Babalola, 2024). For example, co-inoculation of cowpea with *Glomus versiforme* and *Trichoderma harzianum* enhanced seedling growth and resistance to powdery mildew, demonstrating combined fertilization and biocontrol benefits (Kredics *et al.*, 2024). Although field responses may vary due to soil heterogeneity, climatic stress, and inconsistent product quality, the cumulative evidence indicates that locally adapted microbial consortia provide environmentally sustainable and economically viable alternatives to synthetic inputs in Nigerian agriculture (Olaniran *et al.*, 2023; Agbodjato & Babalola, 2024).

CHALLENGES AND OPPORTUNITIES OF MICROBIAL CONSORTIA AS BIOFERTILIZERS AND BIOCONTROL AGENTS IN NIGERIAN AGRICULTURE

Agriculture in Nigeria continues to face declining soil fertility, rising fertilizer costs, and environmental degradation, increasing interest in microbial consortia as sustainable alternatives to chemical inputs. These consortia enhance crop productivity through biological nitrogen fixation, phosphorus solubilization, phytohormone production, and pathogen suppression, thereby improving nutrient use efficiency and plant health. However, adoption remains constrained by technical and institutional barriers. Field performance is often inconsistent due to variations in soil pH, temperature, moisture, and competition with native microbiota, which affect inoculant survival and root colonization (Adesemoye & Egamberdieva, 2013; Ewubare *et al.*, 2023). Short shelf life, heat sensitivity, and poor carrier formulations further reduce product viability during storage and transport (Olanrewaju *et al.*, 2017; Seenivasagan & Babalola, 2021). Unlike chemical fertilizers that supply nutrients immediately, microbial inputs require time to establish, which may discourage farmers seeking rapid yield gains. Limited awareness, weak extension services, and inadequate local production infrastructure also restrict availability and uptake (Babalola, 2010; Ewubare *et al.*, 2023).

Despite these challenges, significant opportunities exist. Microbial consortia lower fertilizer dependence by enhancing nutrient cycling, improve root growth and stress tolerance under drought and nutrient-poor conditions, and suppress soil-borne pathogens through competition and antibiosis. Beneficial genera such as *Trichoderma* and *Pseudomonas* provide effective biological control while reducing pesticide reliance (Seenivasagan & Babalola, 2021). Additionally, reduced chemical use supports soil restoration, lowers greenhouse gas emissions, and creates prospects for local bio-input production using indigenous strains. Advances in formulation technologies and synthetic microbial communities are further improving stability and field reliability, strengthening the case for large-scale adoption across Nigeria agricultural systems.

CONCLUSION AND RECOMMENDATION

Microbial consortia represent a promising frontier in the transition toward biologically based and environmentally sustainable soil fertility and plant health management. Evidence synthesized in this review demonstrates that multispecies inoculants provide broader and more consistent agronomic benefits than single-strain formulations due to functional complementarity, synergistic interactions, and greater ecological resilience in the rhizosphere. Through coordinated mechanisms including biological nitrogen fixation, nutrient solubilization and cycling, phytohormone production, biofilm formation, competitive exclusion, and induced systemic resistance, these consortia enhance nutrient use efficiency, improve soil biological activity, suppress pathogens, and promote crop growth under diverse environmental conditions. Such multifunctionality is particularly relevant for degraded tropical soils where nutrient limitations, low organic matter, and biotic stresses constrain productivity.

Findings from Nigerian agroecosystems further indicate that integrated applications of rhizobia, plant growth-promoting rhizobacteria, and mycorrhizal fungi can substantially increase yields of legumes and cereals while reducing dependence on mineral fertilizers and pesticides. These benefits are economically important for smallholder farmers and environmentally significant for mitigating soil degradation, chemical contamination, and greenhouse gas emissions. Despite these advantages, large-scale adoption remains limited by inconsistent field performance, short shelf life, formulation instability, weak quality assurance systems, limited production infrastructure, and insufficient farmer



awareness. Consequently, the practical effectiveness of microbial consortia depends not only on microbial functionality but equally on formulation science, delivery technologies, regulatory oversight, and knowledge transfer mechanisms.

Future progress therefore requires the systematic development of indigenous, climate-adapted strains and the design of functionally complementary synthetic communities validated through multi-location field trials. Greater emphasis should be placed on improving carrier materials and encapsulation technologies that enhance microbial survival, storage stability, and controlled release in soils. Establishing national standards for inoculant quality, certification, and routine monitoring will be critical to ensure product reliability and build farmer confidence. Strengthened extension services, demonstration trials, and farmer training programs are needed to promote correct application practices and realistic expectations regarding the time-dependent benefits of biological inputs. Local production and commercialization supported through public-private partnerships could further reduce costs, improve accessibility, and stimulate rural bio-input industries. Integrating microbial inoculants into national soil fertility and sustainable agriculture policies, alongside continuous monitoring of soil health and environmental outcomes, will help align these technologies with broader goals of ecological restoration and climate-smart agriculture.

REFERENCES

- Abdulkadir, N. A., Ewusi-Mensah, N., Opuko, A., Yusuf, A. A., Logah, V., Adamu, U. K., Almu H., Sani, A., Muhammad A. A., Hayatu, B., Ahmad, U. B., Lamido, A. K., Aliyu, R. W. & Jibrin, H. J. (2021). Response of groundnut (*Arachis hypogea* L) to rhizobia inoculation, mineral nitrogen and inoculation amendment at the two agro ecological zones of northern Nigerian savannah. *Journal of GSC Advanced Research and Reviews*, **8**(1): 22-27. DOI: 10.30574/gscarr.2021.8.1.0140
- Adesemoye, A. O. & Egamberdieva, D. (2013). Beneficial Effects of Plant Growth-Promoting Rhizobacteria on Improved Crop Production: Prospects for Developing Economies. In: Maheshwari, D., Saraf, M., Aeron, A. (eds) *Bacteria in Agrobiolgy: Crop Productivity*. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-642-37241-4_2
- Agbodjato, N. A., & Babalola, O. O. (2024). Promoting sustainable agriculture by exploiting plant growth-promoting rhizobacteria (PGPR) to improve maize and cowpea crops. *Peer J*, **12**, e16836. <https://doi.org/10.7717/peerj.16836>
- Ashwini, A. (2023). Effect of formulations of microbial consortia on the nutrient content in maize shoot (*Zea mays* L.). *International Journal of Advanced Biochemistry Research*, **7**(2S): 434 – 437. DOI: 10.33545/26174693.2023.v7.i2Sf.250
- Asif, M., Mughal, A. H., Bisma, R., Mehdi, Z., Saima, S., Ajaz, M., Malik, M. A., Masood, A. & Sidique, S. (2018). Application of different strains of biofertilizers for raising quality forest nursery. *International Journal of Current Microbiology and Applied Sciences*, **7**(10), 3680-3686. <https://doi.org/10.20546/ijcmas.2018.710.425>
- Babalola, O. O. (2010). Beneficial bacteria of agricultural importance. *Biotechnology letters*, **32**(11), 1559-1570. <https://doi.org/10.1007/s10529-010-0347-0>
- Bashan, Y., de-Bashan, L. E., Prabhu, S. R., & Hernandez, J. P. (2014). Advances in plant growth-promoting bacterial inoculant technology: formulations and practical perspectives (1998–2013). *Plant and soil*, **378**(1):1 – 33. <https://doi.org/10.1007/s11104-013-1956-x>
- Bebeley, J. F., Kamara, A. Y., Jibrin, J. M., Tofa, A. I., Solomon, R., & Kamai, N. (2024). Effect of combined use of supplementary irrigation, manure and P fertilization on grain yield and profitability of soybean in northern Nigeria. *Heliyon*, **10**(7), e28749. <https://doi.org/10.1016/j.heliyon.2024.e28749>
- Bello, L. A., Saidu, Z. B., & Uzoma, A. O. (2024). Effect of Arbuscular Mycorrhizal Fungi and Nitrogen Sources on Growth and Nutrient Uptake of Maize Plant in Soil of Minna. *Tropical Agroecosystems*, **5**(2): 51 – 55. DOI: <http://doi.org/10.26480/taec.02.2024.51.55>



- Chai, Y. N., Futrell, S., & Schachtman, D. P. (2022). Assessment of Bacterial Inoculant Delivery Methods for Cereal Crops. *Frontiers in microbiology*, 13, 791110. <https://doi.org/10.3389/fmicb.2022.791110>
- Chargrow, (2019). The Biochar Impact: How Biochar Affects Nutrient and Water Retention. Retrieved from <https://char-grow.com/biochar-impact-nutrient-water-retention>
- de Jesús Astacio, L. M., Prabhakara, K. H., Li, Z., Mickalide, H., & Kuehn, S. (2021). Closed microbial communities self-organize to persistently cycle carbon. *Proceedings of the National Academy of Sciences of the United States of America*, 118(45), e2013564118. <https://doi.org/10.1073/pnas.2013564118>
- De Souza, R. S. C., Armanhi, J. S. L., & Arruda, P. (2020). From microbiome to traits: designing synthetic microbial communities for improved crop resiliency. *Frontiers in plant science*, 11, 1179. <https://doi.org/10.3389/fpls.2020.01179>
- Dey, S., Awata, T., Mitsushita, J., Zhang, D., Kasai, T., Matsuura, N., & Katayama, A. (2021). Promotion of biological nitrogen fixation activity of an anaerobic consortium using humin as an extracellular electron mediator. *Scientific reports*, 11(1), 6567. <https://doi.org/10.1038/s41598-021-85955-3>
- Dutta, P., Muthukrishnan, G., Gopalasubramaiah, S. K., Dharmaraj, R., Karuppaiah, A., Loganathan, K., Periyasamy, K., Pillai, M. A., Upamanya, G., Boruah, S., Deb, L., Kumari, A., Mahanta, M., Heisnam, P. & Mishra, A. K. (2022). Plant growth-promoting rhizobacteria (PGPR) and its mechanisms against plant diseases for sustainable agriculture and better productivity. *Biocell*, 46(8), 1843. Doi: 10.32604/biocell.2022.019291
- Egamberdieva, D., Wirth, S. J., Alqarawi, A. A., Abdallah, E. F., & Hashem, A. (2017). Phytohormones and beneficial microbes: essential components for plants to balance stress and fitness. *Frontiers in microbiology*, 8, 2104.
- Etesami, H., & Glick, B. R. (2024). Bacterial indole-3-acetic acid: A key regulator for plant growth, plant-microbe interactions, and agricultural adaptive resilience. *Microbiological research*, 281, 127602. <https://doi.org/10.1016/j.micres.2024.127602>
- Ewubare, P. O., Omokaro, G. O., Ewansiha, O. C., & Edosa, J. O. (2023). Microbial Inoculants in Agriculture: A Microbiological Review from A Nigerian Perspective. *American Journal of Life Science and Innovation*, 2(3), 39 – 43. DOI: <https://doi.org/10.54536/ajlsi.v2i3.2241>
- Feng, D., Li, W., Huang, P., Gu, M., Tang, G., Ding, Y., Cao, G., & Xu, W. (2025). Mechanisms of Microorganisms Alleviating Drought and Salt Stresses in Plants. *Microorganisms*, 13(11), 2565. <https://doi.org/10.3390/microorganisms13112565>
- Fierer, N. (2017). Embracing the unknown: disentangling the complexities of the soil microbiome. *Nature Reviews Microbiology*, 15(10), 579 – 590. <https://doi.org/10.1038/nrmicro.2017.87>
- Gu, S., Yang, T., Shao, Z., Wang, T., Cao, K., Jousset, A., Friman, V., Mallon, C., Mei, X., Wei, Z., Xu, Y., Shen, Q., & Pommier, T. (2020). Siderophore-mediated interactions determine the disease suppressiveness of microbial consortia. *Msystems*, 5(3), 10-1128. <https://doi.org/10.1128/msystems.00811-19>
- Gunal, E. (2025). Biochar-mediated changes in nutrient distribution and leaching patterns: insights from a soil column study. *PeerJ*, 13, e18823. <https://doi.org/10.7717/peerj.18823>
- Herrero, J., Ramírez-Santos, A., Díaz-Santos, E., & Torres-Cortés, G. (2025). Biofertilizers for Enhanced Nitrogen Use Efficiency: Mechanisms, Innovations, and Challenges. *Nitrogen*, 6(4), 111. doi: 10.20944/preprints202510.2490.v1
- Herridge, D. F., Peoples, M. B., & Boddey, R. M. (2008). Global inputs of biological nitrogen fixation in agricultural systems. *Plant and soil*, 311(1): 1-18.
- Herrmann, L., & Lesueur, D. (2013). Challenges of formulation and quality of biofertilizers for successful inoculation. *Applied microbiology and biotechnology*, 97(20): 8859 – 8873. <https://doi.org/10.1007/s00253-013-5228-8>



- Jia, Z., Giehl, R. F., Hartmann, A., Estevez, J. M., Bennett, M. J., & von Wirén, N. (2023). A spatially concerted epidermal auxin signaling framework steers the root hair foraging response under low nitrogen. *Current Biology*, **33**(18): 3926-3941.
- Keerthana, R. S., Patil, C. R., Vithal, N., Gundlur, S. S., & Bhavana, M. (2024). Formulating rhizosphere consortia: Harnessing plant growth-promoting rhizobacteria for enhanced agricultural sustainability. *Journal of Advances in Microbiology Research*; **5**(2): 181 – 184, DOI: <https://doi.org/10.22271/micro.2024.v5.i2c.177>
- Khatri, D. K., Tiwari, D. N., & Bariya, H. S. (2017). Chitinolytic efficacy and secretion of cell wall-degrading enzymes from *Trichoderma* spp. in response to phytopathological fungi. *J. Appl. Biol. Biotechnol.*, **5**(6): 1 – 8. DOI: 10.7324/JABB.2017.50601
- Kredics, L., Büchner, R., Balázs, D., Allaga, H., Kedves, O., Racić, G., Varga, A., Nagy, V. D., Vágvölgyi, C., & Sipos, G. (2024). Recent advances in the use of *Trichoderma*-containing multicomponent microbial inoculants for pathogen control and plant growth promotion. *World journal of microbiology & biotechnology*, **40**(5): 162. <https://doi.org/10.1007/s11274-024-03965-5>
- Kumar, P., Rathee, J., Rani, S., & Suneja, P. (2024). Formulation and effectivity of consortia of potential plant growth promoting endophytic bacteria from chickpea (*Cicer arietinum*). *Journal of Applied Biology & Biotechnology*, **12**(4): 241-247.
- Kumar, S. S., & Mohideen, S. S. (2025). Encapsulation of *L. fermentum* with chitosan-alginate enhances its bioactivity against acrylamide toxicity in *D.mel*. *Scientific reports*, **15**(1): 11324. <https://doi.org/10.1038/s41598-025-95499-5>
- Kumari, R., Pandey, E., Bushra, S., Faizan, S., & Pandey, S. (2024a). Plant growth promoting rhizobacteria (PGPR) induced protection: A plant immunity perspective. *Physiologia Plantarum*, **176**(5), e14495. DOI: 10.1111/ppl.14495
- Kunal, Pranaw, K., Kumawat, K. C., & Meena, V. S. (2023). Plant growth-promoting rhizobacteria (PGPR) and plant hormones: an approach for plant abiotic stress management and sustainable agriculture. *Frontiers in Microbiology*, **14**, 1285756.
- Liu, Y., Gates, A. D., Liu, Z., Duque, Q., Chen, M. Y., Hamilton, C. D., O'Toole, G. A., & Haney, C. H. (2024a). *In vitro* biofilm formation only partially predicts beneficial *Pseudomonas fluorescens* protection against rhizosphere pathogens. *bioRxiv: the preprint server for biology*, 628960. <https://doi.org/10.1101/2024.12.17.628960>
- Liu, Y., Xu, Z., Chen, L., Xun, W., Shu, X., Chen, Y., Sun, X., Wang, Z., Ren, Y., Shen, Q., & Zhang, R. (2024). Root colonization by beneficial rhizobacteria. *FEMS microbiology reviews*, **48**(1). <https://doi.org/10.1093/femsre/fuad066>
- Maciag, T., Krzyzanowska, D. M., Jafra, S., Siwinska, J., & Czajkowski, R. (2020). The Great Five-an artificial bacterial consortium with antagonistic activity towards *Pectobacterium* spp. and *Dickeya* spp.: formulation, shelf life, and the ability to prevent soft rot of potato in storage. *Applied microbiology and biotechnology*, **104**(10): 4547–4561. <https://doi.org/10.1007/s00253-020-10550-x>
- Malusá, E., & Vassilev, N. (2014). A contribution to set a legal framework for biofertilisers. *Applied microbiology and biotechnology*, **98**(15): 6599-6607. doi: 10.1007/s00253-014-5828-y
- Malusà, E., Pinzari, F., & Canfora, L. (2016). Efficacy of biofertilizers: challenges to improve crop production. In *Microbial Inoculants in Sustainable Agricultural Productivity: 2*: 17-40 New Delhi: Springer India.
- Meza, C., Valenzuela, F., Yáñez, M., Cabeza, R. A., Ramos, P., Plaza, A., Carrasco, B., Flores-Castañón, N., Mesquita-Neto, J., Arce-Johnson, P., & Banerjee, A. (2025). Auxin producing plant growth promoting bacteria enhance temperature stress tolerance in Chilean common bean landraces. *Sci Rep* **15**, 30359. <https://doi.org/10.1038/s41598-025-15645-x>
- Niu, B., Wang, W., Yuan, Z., Sederoff, R. R., Sederoff, H., Chiang, V. L., & Borriss, R. (2020). Microbial interactions within multiple-strain biological control agents impact soil-borne plant disease. *Frontiers in Microbiology*, **11**, 585404. <https://doi.org/10.3389/fmicb.2020.585404>



- Nunes, P. S., Lacerda-Junior, G. V., Mascarin, G. M., Guimarães, R. A., Medeiros, F. H., Arthurs, S., & Bettiol, W. (2024). Microbial consortia of biological products: do they have a future? *Biological Control*, 188, 105439. <https://doi.org/10.1016/j.biocontrol.2024.105439>
- Olaniran, A. F., Osemwegie, O., Taiwo, E. A., Okonkwo, C. E., Ojo, O. A., Abalaka, M., Malomo, A. A., Iranloye, Y. M., Akpor, O. B., Bamidele, O. P., & Michael, T. (2023). Application and Acceptability of Microbiomes in the Production Process of Nigerian Indigenous Foods: Drive towards Responsible Production and Consumption. *Preventive nutrition and food science*, 28(2): 108–120. <https://doi.org/10.3746/pnf.2023.28.2.108>
- Olanrewaju, O. S., Glick, B. R., & Babalola, O. O. (2017). Mechanisms of action of plant growth promoting bacteria. *World Journal of Microbiology and Biotechnology*, 33(11): 197. <https://doi.org/10.1007/s11274-017-2364-9>
- Page-Dumroese, D. S., Anderson, N. M., Windell, K. N., Englund, K., & Jump, K. (2016). Development and use of a commercial-scale biochar spreader. *Gen. Tech. Rep. RMRS-GTR-354*. Fort Collins, CO: US Department of Agriculture, Forest Service, Rocky Mountain Research Station. 10:354. <https://doi.org/10.2737/RMRS-GTR-354>
- Philippot, L., Raaijmakers, J. M., Lemanceau, P., & Van Der Putten, W. H. (2013). Going back to the roots: the microbial ecology of the rhizosphere. *Nature reviews microbiology*, 11(11): 789-799. <https://doi.org/10.1038/nrmicro3109>
- Prasad, R., & Chakraborty, D. (2019). Phosphorus basics: understanding phosphorus forms and their cycling in the soil. Extension: Alabama A & M and Auburn Universities. https://www.aces.edu/wp-content/uploads/2019/04/ANR-2535-Phosphorus-Basics_041719L.pdf
- Pretty, J., Benton, T. G., Bharucha, Z. P., Dicks, L. V., Flora, C. B., Godfray, H. C. J., Goulson, D., Hartley S., Lampkin N., Morris C., Pierzynski, G., Prasad, P. V. V., Reganold, J., Rockstrom J., Smith, P., Thorne P. & Wratten, S. (2018). Global assessment of agricultural system redesign for sustainable intensification. *Nature Sustainability*, 1(8): 441-446. <https://www.nature.com/articles/s41893-018-0114-0>
- Qian, X., Chen, L., Sui, Y., Chen, C., Zhang, W., Zhou, J., Dong, W., Jiang, M., Xin, F., & Ochseneither, K. (2020). Biotechnological potential and applications of microbial consortia. *Biotechnology advances*, 40, 107500. <https://doi.org/10.1016/j.biotechadv.2019.107500>
- Rahmati, F. (2020). Microencapsulation of *Lactobacillus acidophilus* and *Lactobacillus plantarum* in Eudragit S100 and alginate chitosan under gastrointestinal and normal conditions. *Applied Nanoscience*, 10(2): 391-399. DOI: 10.1007/s13204-019-01174-3
- Repáč, I., Parobeková, Z., & Sendecký, M. (2017). Reforestation in Slovakia: History, current practice and perspectives. *Reforesta*, (3), 53 – 88. DOI: <https://doi.org/10.21750/REFOR.3.07.31>
- Rivas-García, T., Preciado-Rangel, P., Murillo-Amador, B., Schwentesius-Rindermann, R., Estrada, R. R. G., Hernandez-Vazquez, B., & Torres-Rodríguez, J. A. (2025). Microbial Consortia Used as Biofertilizers and Biocontrol Agents in Agriculture. *Microbial Consortia in Plant Science and Sustainable Agriculture*, 1-14. <https://doi.org/10.1002/9781394325009.ch01>Digital Object Identifier (DOI)
- Saad, M. M., Eida, A. A., & Hirt, H. (2020). Tailoring plant-associated microbial inoculants in agriculture: a roadmap for successful application. *Journal of Experimental Botany*, 71(13): 3878-3901. doi:10.1093/jxb/eraa111
- Saeed, Q., Xiukang, W., Haider, F. U., Kučerik, J., Mumtaz, M. Z., Holatko, J., Naseem, M., Kintl, A., Ejaz, M., Naveed, M., Brtnický, M., & Mustafa, A. (2021). Rhizosphere Bacteria in Plant Growth Promotion, Biocontrol, and Bioremediation of Contaminated Sites: A Comprehensive Review of Effects and Mechanisms. *International journal of molecular sciences*, 22(19): 10529. <https://doi.org/10.3390/ijms221910529>
- Salwan, R., Sharma, M., Sharma, A., & Sharma, V. (2023). Insights into plant beneficial microorganism-triggered induced systemic resistance. *Plant Stress*, 7, 100140. <https://doi.org/10.1016/j.stress.2023.100140>



- Sanginga, N., Okogun, J., Vanlauwe, B., & Dashiell, K. (2002). The contribution of nitrogen by promiscuous soybeans to maize based cropping the moist savanna of Nigeria. *Plant and soil*, **241**(2): 223-231. <https://doi.org/10.1023/A:1016192514568>
- Santoyo, G., Guzmán-Guzmán, P., Parra-Cota, F. I., Santos-Villalobos, S. d. I., Orozco-Mosqueda, M. d. C., & Glick, B. R. (2021). Plant Growth Stimulation by Microbial Consortia. *Agronomy*, **11**(2): 219. <https://doi.org/10.3390/agronomy11020219>
- Seenivasagan, R., & Babalola, O. O. (2021). Utilization of microbial consortia as biofertilizers and biopesticides for the production of feasible agricultural product. *Biology*, **10**(11): 1111. <https://doi.org/10.3390/biology10111111>
- Shahwar, D., Mushtaq, Z., Mushtaq, H., Alqarawi, A. A., Park, Y., Alshahrani, T. S., & Faizan, S. (2023). Role of microbial inoculants as bio fertilizers for improving crop productivity: A review. *Heliyon*, **9**(6). <https://doi.org/10.1016/j.heliyon.2023.e16134>
- Singh, D., Geat, N., Jadon, K. S., Verma, A., Sharma, R., Rajput, L. S., Mahla, H. R., & Kakani, R. K. (2024). Isolation and Characterization of Biocontrol Microbes for Development of Effective Microbial Consortia for Managing *Rhizoctonia bataticola* Root Rot of Cluster Bean Under Hot Arid Climatic Conditions. *Microorganisms*, **12**(11), 2331. <https://doi.org/10.3390/microorganisms12112331>
- Singh, D., Jadon, K. S., Verma, A., Geat, N., Sharma, R., Meena, K. K., & Kakani, R. K. (2025a). Formulations of synergistic microbial consortia for enhanced systemic resistance against Fusarium wilt in cumin. *International microbiology: the official journal of the Spanish Society for Microbiology*, **28**(3): 497–523. <https://doi.org/10.1007/s10123-024-00553-3>
- Singh, M., Jha, S., Pathak, D., & Maisnam, G. (2025). Advancing biofertilizers: the evolution from single-strain formulations to synthetic microbial communities (SynCom) for sustainable agriculture. *Discover Plants*, **2**(1): 226. <https://doi.org/10.1007/s44372-025-00318-w>
- Spaepen, S., & Vanderleyden, J. (2011). Auxin and plant-microbe interactions. *Cold Spring Harbor perspectives in biology*, **3**(4), a001438. <https://doi.org/10.1101/cshperspect.a001438>
- Taiwo, L. B., Adesokun, K. T., Olatoberu, F. T., Oyedele, A. O., Ojo, A. O., & Olayinka, A. A. (2018). Effect of Plant Growth Promoting Rhizobacteria (PGPR) and biochar on soil properties and performance of cowpea (*Vigna unguiculata* (L.) Walp). *Ife Journal of Agriculture*, **30**(3): 56-71. <https://ija.oauiife.edu.ng/index.php/ija/article/view/118>
- Teja, B. S., Gupta, V., Gupta, S. K., Sharma, A., Amin, Z., Mohiddin, F. A., Ashraf, S., Al Tami, M. S., & Hamed, K. E. (2025). Microbial consortia as a biocontrol strategy for bacterial blight in basmati rice. *Revista Argentina de Microbiología*. <https://doi.org/10.1016/j.ram.2025.08.003>
- Tilman, D., Balzer, C., Hill, J., & Befort, B. L. (2011). Global food demand and the sustainable intensification of agriculture. *Proceedings of the national academy of sciences*, **108**(50): 20260-20264. <https://doi.org/10.1073/pnas.1116437108>
- Toju, H., Peay, K. G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., Fukuda, S., Ushio, M., Nakaoka, S., Onoda, Y., Yoshida, K., Schlaeppli, K., Bai, Y., Sugiura, R., Ichihashi, Y., Minamisawa, K. & Kiers, E. T. (2018). Core microbiomes for sustainable agroecosystems. *Nature Plants*, **4**, 247–257. <https://doi.org/10.1038/s41477-018-0139-4>
- Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., & Singh, B. K. (2020). Plant–microbiome interactions: from community assembly to plant health. *Nature reviews microbiology*, **18**(11): 607-621. <https://doi.org/10.1038/s41579-020-0412-1>
- Woo, S. L., & Pepe, O. (2018). Microbial Consortia: Promising Probiotics as Plant Biostimulants for Sustainable Agriculture. *Frontiers in Plant Sciences*, **9**:1801. <https://doi.org/10.3389/fpls.2018.01801>